

Fermion and Anti-Fermion Effective Masses in High Temperature Gauge Theories in CP -Asymmetric Background

A. Erdas

Dipartimento di Scienze Fisiche, Università di Cagliari, 09124 Cagliari, Italy

I.N.F.N. Sezione di Cagliari, 09127 Cagliari, Italy

C. W. Kim and J. A. Lee

Department of Physics and Astronomy

The Johns Hopkins University, Baltimore, MD 21218

Abstract

We calculate the splitting between fermion and anti-fermion effective masses in high temperature gauge theories in the presence of a non-vanishing chemical potential due to the CP -asymmetric fermionic background. In particular we consider the case of left-handed leptons in the $SU(2) \otimes U(1)$ theory when the temperature is above 250 GeV and the gauge symmetry is restored.

erdas@vaxca2.unica.it; cwkim@jhuvms.hcf.jhu.edu; lee@dirac.pha.jhu.edu

I. INTRODUCTION

The behavior of gauge theories at finite temperature and density has been extensively investigated in the literature. In particular, the modifications to the fermion dispersion relation due to high-temperature effects for a chirally invariant gauge theory have been studied by Weldon [1] who demonstrated that right-handed and left-handed leptons in the $SU(2) \otimes U(1)$ model acquire different effective masses. The properties of neutrinos at low temperature and density have also been studied by many authors [2–4].

In this article we examine the properties of chirally invariant gauge theories at high temperature, in particular in the presence of a non-vanishing chemical potential due to a CP-asymmetric fermionic background; one of these theories can be, for example, the Standard Electroweak Model at temperatures for which the $SU(2) \otimes U(1)$ symmetry is restored. The symmetry restoration that occurs in the Standard Electroweak Model at high enough temperature has been studied extensively [5]: above $T = 250\text{GeV}$ the Higgs vacuum expectation value vanishes and so do the masses of the gauge bosons, scalars and the fermions.

The interaction of a particle with the heat bath changes the dispersion relation from the tree level relation. Weldon [1] calculated this temperature correction for chirally symmetric gauge theories in a CP-symmetric medium at high temperature. The purpose of this paper is to extend the Weldon's work to include the case of a non-vanishing chemical potential for the fermions. In the scenario in which the baryogenesis takes place right after the GUTS phase transition and $B - L$ conservation is assumed, the number of particles in the $SU(2) \otimes U(1)$ symmetric phase of the early universe could be slightly larger than the number of antiparticles, implying non-vanishing chemical potentials and leading to the current particle-antiparticle asymmetry [6,7].

To compute the effective fermion masses under the above condition, we need to evaluate the fermion self-energy [1]

$$\Sigma(K) = -a\gamma_\mu K^\mu - b\gamma_\mu u^\mu, \quad (1.1)$$

where u^α is the four-velocity of the heat bath, K^α is the four-momentum of a fermion under consideration and a and b are Lorentz-invariant functions that depend on the Lorentz-scalars defined by [1]

$$\omega \equiv K^\alpha u_\alpha, \quad k \equiv [(K^\alpha u_\alpha)^2 - K^2]^{1/2}. \quad (1.2)$$

We then find the poles of the fermion propagator

$$S(K) = [(1 + a)\gamma_\mu K^\mu + b\gamma_\mu u^\mu]^{-1}. \quad (1.3)$$

In Section II we compute the fermion self-energy in the presence of a non-vanishing chemical potential. Using the results of Section II, we find, in Section III, an integral equation for the effective mass in the presence of the chemical potential $\mu \ll T$ and use this integral equation to calculate the mass-splitting between fermions and anti-fermions. Finally in Section IV we discuss some implications of this mass splitting for left-handed leptons of the $SU(2) \otimes U(1)$ model for $T > 250\text{GeV}$ and in a CP-asymmetric medium.

II. COMPUTATION OF THE SELF-ENERGY

Let us consider a non-abelian gauge field theory in which the fermions, gauge bosons and scalars do not have masses [1]. We are interested in calculating the fermion self-energy at the one-loop level at finite temperature in the presence of a CP-asymmetric fermionic background. We assume the chemical potentials μ_{f_i} of the different fermion species f_i in the theory to be the same

$$\mu_{f_1} = \mu_{f_2} = \cdots = \mu_{f_n} = \mu. \quad (2.1)$$

We use the real-time formulation of finite-temperature field theory and perform the calculation in the Feynman gauge. The three relevant Feynman diagrams are shown in Fig. 1.

At tree level the free-particle (massless) propagators for fermion, gauge boson and scalar in the Feynman gauge are given by

$$S(p) = \gamma_\mu p^\mu \left[\frac{1}{p^2 + i\epsilon} + i\Gamma_f(p) \right], \quad (2.2)$$

$$D_{\mu\nu}(p) = -\eta_{\mu\nu} \left[\frac{1}{p^2 + i\epsilon} - i\Gamma_b(p) \right], \quad (2.3)$$

$$D(p) = \frac{1}{p^2 + i\epsilon} - i\Gamma_b(p), \quad (2.4)$$

where $\Gamma_f(p)$ and $\Gamma_b(p)$ are defined as

$$\Gamma_b(p) \equiv 2\pi\delta(p^2)n_b(p) \quad (2.5)$$

$$\Gamma_f(p) \equiv 2\pi\delta(p^2)n_f(p) \quad (2.6)$$

with

$$n_b(p) = \left[e^{|p \cdot u|/T} - 1 \right]^{-1} \quad (2.7)$$

$$n_f(p) = \theta(p \cdot u)n_f^-(p) + \theta(-p \cdot u)n_f^+(p). \quad (2.8)$$

In Eq. (2.8) the two functions $n_f^\mp(p)$ (not to be confused with the fermion number density N_f that will be introduced later) refer, respectively, to fermions and anti-fermions. These differ from $n_f(p)$ in the Weldon's paper in that they contain the chemical potential μ_f of the fermion f

$$n_f^\mp(p) = \left[e^{(|p \cdot u| \mp \mu_f)/T} + 1 \right]^{-1}. \quad (2.9)$$

The chemical potential μ_f is related to the number densities of fermions N_f and anti-fermions $N_{\bar{f}}$, and is given, in the case of $\mu_f \ll T$, by

$$N_f - N_{\bar{f}} \equiv \int \frac{d^3p}{(2\pi)^3} \left[n_f^-(p) - n_f^+(p) \right] \simeq \frac{\mu_f T^2}{6}. \quad (2.10)$$

We first consider the tadpole diagrams: if all the particles in the theory are massless at tree-level and if $\mu_{f_1} = \mu_{f_2} = \dots = \mu_{f_n}$ then the sum of all the relevant tadpole diagrams will be proportional to

$$\sum_m L_{mm}^A = 0, \quad (2.11)$$

where L_{mm}^A are the representation matrices of the group generators, A runs over the generators and the two lower indices run over the states of the fermion representation [1]. Therefore we can see that the contributions from all the tadpole diagrams add up to zero.

Since the calculations of the two diagrams in Fig. 1 are very similar, we only will show some details of the calculation of the gauge boson contribution to the one-loop fermion self-energy. Taking $\mu_{f_1} = \mu_{f_2} = \dots = \mu_{f_n} = \mu$ the sum of the gauge bosons contribution is given by

$$\Sigma(K) = ig^2 C(R) \int \frac{d^4 p}{(2\pi)^4} D_{\mu\nu}(p) \gamma^\mu S(p+K) \gamma^\nu, \quad (2.12)$$

where g is the coupling constant and $C(R)$ is the quadratic Casimir invariant of the gauge group representation. We are only interested in the real part [1] of the finite temperature corrections Σ' to the fermion self-energy

$$\Sigma' = \Sigma - \Sigma(T=0). \quad (2.13)$$

Substituting the expressions for the propagators Eqs. (2.2) and (2.3) into Eq. (2.13), we obtain

$$Re\Sigma' = 2g^2 C(R) \int \frac{d^4 p}{(2\pi)^4} \left[(\gamma_\mu p^\mu + \gamma_\mu K^\mu) \frac{\Gamma_b(p)}{(p+K)^2} - (\gamma_\mu p^\mu + \gamma_\mu K^\mu) \frac{\Gamma_f(p+K)}{p^2} \right]. \quad (2.14)$$

Since Eq. (2.14) is manifestly covariant, it must be a linear combination of $\gamma_\mu K^\mu$ and $\gamma_\mu u^\mu$. Following the procedure of the Weldon's paper [1], we obtain, after performing the integration over p_0 and over the angles,

$$\begin{aligned} \frac{1}{4} Tr(\gamma_\mu K^\mu Re\Sigma') &= g^2 C(R) \int \frac{dp}{8\pi^2} \left\{ \left[4p - \frac{K^2}{2k} L_1(p) \right] n_b(p) + \right. \\ &\quad + \left[2p + \frac{K^2}{2k} \left(\ln \left[\frac{p+\omega_+}{p+\omega_-} \right] - \ln \left[\frac{\omega_+}{\omega_-} \right] \right) \right] n_f^-(p) + \\ &\quad \left. + \left[2p + \frac{K^2}{2k} \left(-\ln \left[\frac{p-\omega_+}{p-\omega_-} \right] + \ln \left[\frac{\omega_+}{\omega_-} \right] \right) \right] n_f^+(p) \right\}, \end{aligned} \quad (2.15)$$

$$\begin{aligned} \frac{1}{4} Tr(\gamma_\mu u^\mu Re\Sigma') &= \frac{g^2 C(R)}{k} \int \frac{dp}{8\pi^2} \left[\left(2p \ln \left[\frac{\omega_+}{\omega_-} \right] - p L_2(p) - \omega L_1(p) \right) n_b(p) \right. \\ &\quad \left. - p \left(\ln \left[\frac{p+\omega_+}{p+\omega_-} \right] - \ln \left[\frac{\omega_+}{\omega_-} \right] \right) n_f^-(p) - p \left(\ln \left[\frac{p-\omega_+}{p-\omega_-} \right] - \ln \left[\frac{\omega_+}{\omega_-} \right] \right) n_f^+(p) \right], \end{aligned} \quad (2.16)$$

where

$$\omega_\pm = \frac{1}{2}(\omega \pm k)$$

$$L_1(p) = \ln \left[\frac{p + \omega_+}{p + \omega_-} \right] - \ln \left[\frac{p - \omega_+}{p - \omega_-} \right]$$

$$L_2(p) = \ln \left[\frac{p + \omega_+}{p + \omega_-} \right] + \ln \left[\frac{p - \omega_+}{p - \omega_-} \right] .$$

Here all logarithms are to be understood in the principal-value sense.

III. CALCULATION OF THE EFFECTIVE MASS SPLITTING

The two quantities in Eqs. (2.15) and (2.16) can be written in terms of the Lorentz-invariants a and b

$$\frac{1}{4} \text{Tr}(\gamma_\mu K^\mu \text{Re}\Sigma') = -aK^2 - bK \cdot u = -a(\omega^2 - k^2) - b\omega \quad (3.1)$$

$$\frac{1}{4} \text{Tr}(\gamma_\mu u^\mu \text{Re}\Sigma') = -aK \cdot u - b = -a\omega - b , \quad (3.2)$$

from which we obtain

$$ak^2 = \frac{1}{4} \text{Tr}(\gamma_\mu K^\mu \text{Re}\Sigma') - \omega \frac{1}{4} \text{Tr}(\gamma_\mu u^\mu \text{Re}\Sigma') \quad (3.3)$$

$$bk^2 = -\omega \frac{1}{4} \text{Tr}(\gamma_\mu K^\mu \text{Re}\Sigma') + (\omega^2 - k^2) \frac{1}{4} \text{Tr}(\gamma_\mu u^\mu \text{Re}\Sigma') . \quad (3.4)$$

The poles in the propagator occur, as can be seen from Eq. (1.3), when ω and k are such as to produce a zero in the following Lorentz-invariant function

$$D = (1 + a)^2 K^2 + b^2 + 2(1 + a)bK \cdot u . \quad (3.5)$$

The positive-energy root occurs at

$$\omega(1 + a) + b = k(1 + a) . \quad (3.6)$$

We want to find the solution of this equation for $k = 0$. We will call M the value of ω that satisfies Eq. (3.6) for $k = 0$:

$$\omega(k = 0) = M \quad (3.7)$$

and M can be interpreted as an effective fermion mass.

For small k , the two expressions $\frac{1}{4}Tr(\gamma_\mu K^\mu Re\Sigma')$ and $\frac{1}{4}Tr(\gamma_\mu u^\mu Re\Sigma')$ are of the form

$$\frac{1}{4}Tr(\gamma_\mu K^\mu Re\Sigma') = h_0 + h_1 k^2 + h_2 k^4 + \dots \quad (3.9)$$

$$\frac{1}{4}Tr(\gamma_\mu u^\mu Re\Sigma') = g_0 + g_1 k^2 + g_2 k^4 + \dots , \quad (3.10)$$

where the h_i and g_i are functions of ω and T alone.

We now want to use the expressions given by Eqs. (3.9) and (3.10) to find the expression of a and b for small k . Inserting Eq. (3.9) and (3.10) into (3.3) and (3.4), we get

$$ak^2 = h_0 + h_1 k^2 - \omega g_0 - \omega g_1 k^2 + \dots \quad (3.11)$$

$$bk^2 = -\omega h_0 - \omega h_1 k^2 + (\omega^2 - k^2)(g_0 + g_1 k^2) + \dots . \quad (3.12)$$

Since we have

$$h_0 - \omega g_0 = 0 , \quad (3.13)$$

we obtain

$$a \simeq h_1 - \omega g_1 \quad (3.14)$$

$$b \simeq -\omega h_1 + \omega^2 g_1 - g_0 . \quad (3.15)$$

Using now Eqs. (3.14) and (3.15), we can rewrite Eq. (3.6) for small k as

$$\omega(1 + h_1 - \omega g_1) - \omega h_1 + \omega^2 g_1 - g_0 = k(1 + h_1 - \omega g_1) \quad (3.16)$$

which, after simplification, becomes

$$\omega - k - g_0 = k(h_1 - \omega g_1) \quad (3.17)$$

and for $k = 0$ gives

$$M = g_0 \quad (3.18)$$

and therefore

$$M^2 = h_0 . \quad (3.19)$$

The quantity h_0 is given, from Eq. (3.9), by

$$h_0 = \lim_{k \rightarrow 0} \frac{1}{4}Tr(\gamma_\mu K^\mu Re\Sigma') . \quad (3.20)$$

After a lengthy but straightforward algebra, Eq.(3.20) yields

$$h_0 = \frac{g^2 C(R)}{8\pi^2} \left\{ 2 \left(\frac{T^2 \pi^2}{3} + \frac{T^2 \pi^2}{6} + \frac{\mu^2}{2} \right) - M\mu - M^2 \int dp \left(\frac{1}{p+M} - \frac{1}{M-p} \right) n_b(p) \right. \\ \left. + M^2 \int dp \left[\frac{1}{2p+M} \left(n_f^-(p) - \frac{1}{e^{p/T} + 1} \right) - \frac{1}{M-2p} \left(n_f^+(p) - \frac{1}{e^{p/T} + 1} \right) \right] \right\}. \quad (3.21)$$

Therefore, in order to find the effective fermion mass, we have to solve the following equation for M

$$M^2 = M_0^2 \left\{ 1 + \left(\frac{\mu}{\pi T} \right)^2 - \frac{M\mu}{\pi^2 T^2} - \frac{M^2}{\pi^2 T^2} \int_0^\infty dp \left(\frac{1}{p + \frac{M}{T}} - \frac{1}{\frac{M}{T} - p} \right) \frac{1}{e^p - 1} \right. \\ \left. + \frac{M^2}{\pi^2 T^2} \int dp \left[\frac{1}{2p+M} \left(\frac{1}{e^{(p-\mu)/T} + 1} - \frac{1}{e^{p/T} + 1} \right) - \frac{1}{M-2p} \left(\frac{1}{e^{(p+\mu)/T} + 1} - \frac{1}{e^{p/T} + 1} \right) \right] \right\}, \quad (3.22)$$

where

$$M_0^2 = g^2 \frac{C(R)}{8} T^2 \quad (3.23)$$

is the value of the fermion effective mass in the absence of a chemical potential [1]. When we also add the scalar contributions to the fermion self-energy, M_0^2 is modified to

$$M_0^2 = g^2 \frac{C(R)}{8} T^2 + \frac{C'|f|^2}{16} T^2, \quad (3.24)$$

where $|f|$ is the Yukawa coupling constant of the fermions and C' is a numerical constant that depends on the fermion representations [1]. Equation (3.22) reduces to the result in Ref.[1] in the limit $\mu \rightarrow 0$, as it should.

We now solve Eq.(3.22) for M in the limit $\mu \ll T$. In this limit the second line of Eq. (3.22) becomes

$$\frac{M^2}{\pi^2 T^2} \int dp \left[\frac{1}{2p+M} \left(\frac{1}{e^{(p-\mu)/T} + 1} - \frac{1}{e^{p/T} + 1} \right) - \frac{1}{M-2p} \left(\frac{1}{e^{(p+\mu)/T} + 1} - \frac{1}{e^{p/T} + 1} \right) \right] \simeq \\ \frac{M^2}{\pi^2 T^2} \frac{\mu}{T} \int dp \left(\frac{1}{2p+M} + \frac{1}{M-2p} \right) \frac{1}{e^{p/T} + 1} \left(1 - \frac{1}{e^{p/T} + 1} \right). \quad (3.25)$$

Defining

$$M \equiv M_0 + x, \quad (3.26)$$

with

$$x \ll M_0, \quad (3.27)$$

and neglecting higher order terms (such as the one in Eq. (3.25)) and the contribution from the Yukawa couplings (i.e. $|f|^2 \ll g^2$), we obtain

$$x \simeq \frac{M_0}{2} \left[-g \sqrt{\frac{C(R)}{8\pi^4}} \frac{\mu}{T} - g^2 \frac{C(R)}{8\pi^2} \int_0^\infty dp \left(\frac{1}{p + g \sqrt{\frac{C(R)}{8}}} - \frac{1}{g \sqrt{\frac{C(R)}{8}} - p} \right) \frac{1}{e^p - 1} \right]. \quad (3.28)$$

In general we have

$$\frac{\mu}{T} \sim g^n,$$

where the integer n will depend on the specific gauge theory and temperature range of interest, therefore, using also Eq. (3.23), we can see that the value of the fermion effective mass is given, up to order g^{n+1} , by

$$M = M_0(1 + c_1 g + c_2 g^2 + \dots + c_n g^n + c_{n+1} g^{n+1} - \frac{g}{2} \sqrt{\frac{C(R)}{8}} \frac{\mu}{T} + \dots). \quad (3.29)$$

In order to find the values of the c_i (which are independent of μ and T) we would need to solve numerically the integral of Eq. (3.28) and to include higher order corrections to the fermion self-energy (two-loops etc.). The effective mass $\overline{M}$ of the antifermions can be obtained by replacing μ by $-\mu$ in M . Therefore, we find

$$\Delta M \equiv \overline{M} - M = g^2 \frac{C(R)}{8\pi^2} \mu = \frac{3}{4\pi^2} \frac{g^2 C(R)}{T^2} (N_f - N_{\bar{f}}). \quad (3.30)$$

IV. DISCUSSION

We have calculated the effective mass splitting between fermions and anti-fermions in the CP-asymmetric background at high temperature to be $\Delta M = \overline{M} - M = g^2 \frac{C(R)}{8\pi^2} \mu$. Now, let us consider the case of $SU(2) \otimes U(1)$ at $T > 250\text{GeV}$ (when the gauge symmetry is still unbroken) and with a CP -asymmetric fermionic background. Concentrating on the lepton sector of the first generation, we can take the chemical potential μ_{e_L} of the left-handed

electrons, μ_{e_R} of the right-handed electrons and the chemical potential μ_ν of the neutrinos to be the same

$$\mu_{e_L} = \mu_{e_R} = \mu_\nu = \mu . \quad (4.1)$$

(As pointed out in Ref. [8], if one considers only one generation of weak doublet, due to the effect of electroweak fermion number violation $N_f - N_{\bar{f}}$ would become zero. However, with more than one doublet, the difference between any two fermion numbers is conserved and the equilibrium number density $N_f - N_{\bar{f}}$ at a given temperature for a given generation will not be zero.)

In this case the mass splitting ΔM between a left-handed anti-lepton and its anti-particle becomes

$$\Delta M = \frac{g'^2 + 3g^2}{32\pi^2} \mu , \quad (4.2)$$

where g' is the coupling constant of the $U(1)$ field and g is the coupling constant of the $SU(2)$ gauge fields. In deriving Eq.(4.2) we have used $C(R) = \frac{3}{4}$ and 1 for $SU(2)$ and $U(1)$ gauge groups, respectively, and the fact that the left-handed fermions couple to the $U(1)$ gauge field with the strength $g'/2$.

Using Eq. (2.10) and

$$N_{e_L} - N_{\bar{e}_L} \simeq N_\nu - N_{\bar{\nu}} = N_f - N_{\bar{f}}$$

we can write

$$\Delta M = \frac{3}{16\pi^2} (g'^2 + 3g^2) \frac{N_f - N_{\bar{f}}}{T^2} . \quad (4.3)$$

Taking, based on $B - L$ conservation,

$$\frac{N_f - N_{\bar{f}}}{N_\gamma} \simeq \frac{N_B - N_{\bar{B}}}{N_\gamma} \sim 10^{-9}, \quad (4.4)$$

we see that

$$N_f - N_{\bar{f}} \sim 10^{-9} T^3 . \quad (4.5)$$

Therefore, it appears that the value of ΔM we obtained in Eq. (4.3) is too small to play a significant role in the evolution of the $N_f - N_{\bar{f}}$ and consequently $N_B - N_{\bar{B}}$ in the early

Universe. Possible effects of Eq. (4.3) on neutrino oscillations in the early Universe will be given elsewhere.

ACKNOWLEDGMENTS

One of us (A.E.) wishes to thank the High Energy Theory Group of the Johns Hopkins University for the hospitality extended to him during his several visits. This work was supported in part by the National Science Foundation.

REFERENCES

- [1] H. A. Weldon, Phys. Rev. D 26, 2789 (1982)
- [2] D. Notzold, G. Raffelt, Nucl. Phys. B307, 924 (1988)
- [3] S. S. Masood, Preprint CERN-TH.6622/92 (1992)
- [4] J. F. Nieves, Phys. Rev. D 40, 866 (1989)
- [5] M. Sher, Phys. Rep. 179, 273 (1989)
- [6] M. Dine, Preprint SCIPP 92/91 (1992)
- [7] A. G. Cohen et al., to appear in Ann. Rev. of Nucl. and Part. Sci., 43 (1993)
- [8] S. M. Barr, R. S. Chivukula and E. Farhi, Phys. Lett. B241, 387 (1990).

Figure Caption

FIG. 1. The 3 diagrams relevant to the one-loop self-energy calculation; (a) the tadpole, (b) the gauge boson contribution and (c) the scalar contribution.